

Comparison of Combination of Sclerotherapy and Tiersch's Procedure with Perineal Rectopexy with Mesh in Children

Reza Ebrahimi¹, Baharah Ahmadi¹, Mehrdad Hosseinpour^{1*}

¹Imam Hossein Children Hospital, Isfahan University of Medical Sciences, Isfahan, Iran

***Address for Corresponder:** Dr. Mehrdad Hosseinpour, Imam Hossein Children Hospital, Isfahan University of Medical Sciences, Isfahan, IRAN (email: meh_hosseinpour@yahoo.com)

How to cite this article:

Ebrahimi R, Ahmadi B, Hosseinpour M. Comparison of Combination of Sclerotherapy and Tiersch's Procedure with Perineal Rectopexy with Mesh in Children. *Iranian Journal of Pediatric Surgery* 2020; 6 (2): 74-80.

DOI: <https://doi.org/10.22037/irjps.v6i2.30061>

Abstract

Introduction: Perineal rectopexy is a method for surgical treatment of rectal prolapse in children, but it is accomplished using different techniques. We investigated the combination therapy of Tiersch's operation and sclerotherapy and compared its outcome with mesh rectopexy with talc.

Materials and Methods: A total of 80 children with rectal prolapse were selected and divided in two groups. In the control group rectopexy was performed by perineal mesh rectopexy with talc. In the case group, injection of 30% saline and, Tiersch's operation were performed. The parameters used to compare the success of interventions were improvement of incontinence and constipation and a low rate of recurrence.

Result: Our results showed that the rate of improvement in constipation and incontinence was not significantly different in the two groups. The recurrence rate was 1.6 % in both groups in one year of follow up. Fifteen patients (37.5%) in the control group had severe anxiety and pain during the extraction of mesh. The length of hospitalization was 6.2 ± 0.94 in the controls and 0.86 ± 0.63 days in case group.

Conclusion: Our study suggests the combination perineal therapy (T+S) as an alternative approach for children who have limitations regarding talc rectopexy or the high hospital charges related to this approach.

Keywords

- Rectal Prolapse
- Sclerotherapy
- Rectopexy

Introduction

Rectal prolapse in young children is a common disease which is treated conservatively. Although most cases of prolapse is mucosal, occasionally it is full thickness and therefore does not respond to nonsurgical treatments¹⁻². Many treatment options are available such as injection of sclerosant, abdominal or perineal surgical operations³⁻⁵ and they have different success rates⁶⁻⁸, the optimal management of this disease is still under debate.⁹

Here, we compared perineal mesh rectopexy with combination therapy of Tiersch's operation plus sclerotherapy (T+S) and discussed the success rate of these techniques and their related complications.

Materials and Methods

In a randomized controlled trial carried out in three large children hospitals in Iran from 2006 to April 2020; the effect of therapeutic interventions on two groups of children admitted for elective surgery of rectal prolapse; cases and controls were examined. Children were divided into 2 groups, children with odd hospital record numbers were assigned to the case group and those with even numbers were allocated to the control group.

Inclusion criteria were as follows: 1- no response to conservative management (laxatives, high fiber diet, regular prompt defecation from a sitting position) 2- failure of conservative management as shown by recurrence of the prolapse and 3- Full thickness prolapse.

Exclusion criteria were as follows: children with medical diseases or a positive family history of

pulmonary disease, lung cancer, skin or ovarian cancers.¹⁰⁻¹¹

The study was approved by our hospital ethics committee and written consent was obtained.

Prolapse diagnosis was made clinically and we didn't use defecography or MRI in this study. On pre-operative evaluation, demographic data, prolapse duration, clinical findings (Protrusion, constipation, diarrhea, bleeding and incontinency), and duration of conservative management were collected.

Techniques:

With general anesthesia, in lithotomy position, after initial rectal examination, proctoscopy was performed. If rectum was loaded with hard stool, it was evacuated.

In the controls, with a 180° semi-circular incision of the posterior anal verge (between coccyx and anus), the presacral space was dissected up to the sacral promontory. A 20 cm mesh which was soaked in asbestos-free talc (Bryan Corp, Woburn, MA, USA),¹³⁻¹⁴ was inserted in the pre sacral space¹² with the ending hanging out of the incision. It was extracted gradually (6 cm /day) from the 5th day of surgery.

In the case group, the anus was dilated using a loaner retractor. A long 23- gauge needle was placed into the sub-mucosal plane approximately 4 cm from the anal verge and 2 ml of 30 percent solution was injected into each of three quadrants (3, 6 and 9 o'clock).¹⁵ Sufficient sclerosant injection was achieved by bulging or blanching of the injection

site. Then, with two small incisions at 12 and 6 o'clock of the anal verge, a length of absorbable (0 caliber Vicryl) suture was threaded from the posterior incision to the anterior incision around the anus, just deep to the external sphincter muscle (ring formation). The suture was pulled and tied on a Hegar's dilator. Absorbable sutures were used to close the two incisions.¹⁶

On postoperative assessment, hospitalization length and post operative complications (infection, bleeding) were recorded. A one year follow up was carried out for all patients. The main comparative variables were improvement of incontinence and constipation and a low rate of recurrence.

For data analysis we used the SPSS software. Data regarding quantitative variables was expressed as mean \pm SD and Mann-Whitney test was used for

comparing the mean scores. Qualitative data and there relation to different operative techniques was assessed using chi-square and Fisher's exact test. P values less than 0.05 was considered significant

Results

All 80 children (40 cases and 40 controls) were evaluated. The mean age was 5.1 ± 0.09 (years) in controls and 4.82 ± 0.63 in cases ($P > 0.005$). There was no significant difference in the sex of patients (male/female: 22/18 in controls, 26/14 in cases). Duration of prolapse prior to treatment was 14.2 ± 4.2 months in controls and 14.6 ± 3.7 months in cases; which was not significant. Duration of conservative management was 12.2 ± 2.9 months in controls and 14.3 ± 1.9 months in cases (also not significant). Clinical findings of the patients are demonstrated in **Table 1**.

Table 1: Clinical findings of groups

P	Control [n (%)]	Case [n (%)]
Protrusion NS	40(100)	40(100)
Constipation NS	21(52.5)	26(65)
Diarrhea NS	2(5)	3(7.5)
Bleeding NS	1(2.5)	0(0)
Incontinence NS	5(12.5)	3(7.5)

Postoperative complication occurred only in 2 patients of the control group: 1 postoperative infection (2.5%) and 1 bleeding (2.5%) which were not significantly different.

15 patients (37.5%) in the control group had severe

anxiety and pain during the extraction of mesh. The sweat test was positive in two patients.

Table 2 shows the success rate of different techniques.

Table 2: Success rate in children after surgery

Parameter P	Group (n)	Success rate [n (%)]
Constipation NS	Control (27)	26(96.6)
	Case (21)	20(96.6)
Incontinence NS	Control (3)	2 (66)
	Case (5)	5(100)
Recurrence NS	Control (40)	1 (1.6)
	Case (40)	1 (1.6)

The results showed that there was a higher but not significant resolution rate for incontinency in the case group.

Duration of Hospitalization was 6.2 ± 0.94 days in the controls and 0.86 ± 0.63 days in case the group (P: not significant).

Discussion

Although surgical severe rectal prolapse is commonly a disease of old age;¹⁷⁻¹⁸ in children conservative management is the main treatment. However in some cases, surgical management is mandatory. Despite the progresses in modern surgery (laparoscopic rectopexy and injection¹⁹), choosing the surgical procedure for management of rectal prolapse is still a dilemma.

The basic underlying mechanism leading to rectal prolapse in children is stretching and weakening of the levator muscle complex surrounding the rectal wall. The most common predisposing causes are disorders of sacral nerve root innervations, chronic malnutrition, acquired muscular weakness associated with straining at defecation and idiopathic

prolapse which is the most common cause in our series.

As previously described by Nazem et al.,¹⁶ perineal mesh rectopexy is an alternative approach for abdominal operations in children. Sterile talc powder as a sclerosing agent had acceptable results with low rate of complication in treatment of rectal prolapse in children. Talc stimulates an inflammatory reaction in the perineal region and results in better adherence of the rectum to the sacrum.

However, we have three major limitations in mesh rectopexy with talk. Firstly, although the US Food and Drug Administration (FDA) considered the non-asbestiform talc as a safe agent, but there are some studies showing a preliminary link between talk and some types of cancers.¹⁰⁻¹¹ Secondly, some patients experienced a high degree of anxiety and pain during the extraction of mesh. Thirdly, the length of hospitalization and patient payment is also considerable in mesh rectopexy. Therefore, we used a combination therapy of T+S as an alternative approach for surgical treatment of rectal prolapse.

Our findings showed that length of hospitalization is shorter in T+S method and most patients can be discharged on the first day of operation. The rate of postoperative constipation was low in both methods, because in these approaches, the parasympathetic nerve fibers of lateral ligaments of rectum remain intact.²⁰

Other findings of this study showed improvement of incontinency in patients especially in the T+S group. Incontinence in rectal prolapse can be related to pudendal nerve latency, external sphincter denervation,²¹ persistent recto- anal inhibition, low rectal sensory thresholds, adaptation to distension²² and hypermotility of sigmoid and descending colon.²³

Although it seems that rectopexy by any method improves internal sphincter function, our findings show that the addition of narrowing of the anal canal by sclerotherapy had a better result.

On the other hand, prolonged follow up of cases treated by sclerotherapy shows a recurrence rate of 10% in two years²⁴ but in combination therapy, the results are comparable to other surgical techniques

such as mesh rectopexy.²⁵⁻²⁶

Conclusion

Our study suggests the combination perineal therapy (T+S) as an alternative approach for children who have limitations regarding talc rectopexy or the high hospital charges related to this approach.

Ethical Consideration

In this study, principles of ethical and professional conduct of our hospital and university have been followed and Ethic number of Imam Hossein Hospital is 100125.

Acknowledgements

Not applicable

Funding/Support

We have no sources for funding in this study.

Conflict of interests

There is no conflict of interest.

References

1. Antao B, Bradley V, Roberts JP, et al: Management of rectal prolapse in children. *Dis Colon Rectum* 2005;48(8):1620-5.
2. Madiba TE1, Baig MK, Wexner SD: Surgical management of rectal prolapse. *Arch Surg* 2005; 140 (1):63-73.
3. Groff DB, Nagaraj HS: Rectal prolapse in infants and children. *Am JSurg* 1990; 160 (5):531-2.
4. Zempsky WT, Rosenstein BJ: The cause of rectal prolapse in children. *Am J Dis Child* 1988; 142(3):338-9.

5. Hetzer FH, Bieler A, Hahnloser D: Outcome and cost analysis of sacral nerve stimulation for fecal incontinence. *Br J Surg* 2006; 93 (11): 1411 – 1417
6. Qaradaghy SH, Hawramy TA, Nore BF, et al: Longitudinal plication--a surgical strategy for complete rectal prolapse management. *BMC Surg* 2014; 14:17.
7. Hata F, Nishimori H, Ikeda S, et al: A simple and safe procedure to repair rectal prolapse perineally using stapling devices. *Case Rep Gastroenterol* 2014; 8(1):39-4.
8. Bergamaschi R, Corman ML: The rectal prolapse dilemma: abundance of adjectives. *Cir Esp* 2014; 92(3):147-8.
9. Calkins C, Oldham K: Acquired anorectal disorders, in: Holcomb GW, Murphy JP, Ostlie DJ, eds: *Ashcraft Pediatric Surgery*. Philadelphia: Elsevier Saunders; 2014, 525 – 529.
10. Hollinger MA: Pulmonary toxicity of inhaled and intravenous talc. *Toxicol Lett* 1990; 52 (2): 121 – 127.
11. Harlow BL, Cramer DW, Bell DA et al: Perineal exposure to talc and ovarian cancer risk. *Obstetrics and Gynecology* 1992; 80 (1): 19 – 26.
12. Nazem M, Hosseinpour M, Farhadi M: Perineal Mesh Rectopexy with Sterile Talc in Children with Rectal Prolapse. *Eur J Pediatr Surg* 2010; 20: 321– 324.
13. Yim AP, Chan AT, Lee TW: Thorascopic talc insufflation versus talc slurry for symptomatic malignant pleural effusion. *Ann Thorac Surg* 1996; 62 : 1655 – 1658.
14. Ong KC, Indumathi V, Raghuram J, et al: A comparative study of pleurodesis using talc slurry and bleomycin in the management of malignant pleural effusions. *Respirology* 2000; 5 (2): 99 – 103.
15. Abes M, Sarihan H: Injection sclerotherapy of rectal prolapse in children with 15 percent saline solution. *Eur J Pediatr Surg* 2004; 14:100–2.
16. Coppi P: Rectal prolapse, in Spitz L, Coran AG, (eds): *Operative pediatric surgery*. Abingdon: Taylor and Francis, 2013, 608-614.
17. Bhandarkar DS: Laparoscopic rectopexy for complete rectal prolapse: mesh, no mesh or a ventral mesh? *J Minim Access Surg* 2014; 10(1):1-3.
18. Lee JL, Yang SS, Park IJ, et al: Comparison of abdominal and perineal procedures for complete rectal prolapse: an analysis of 104 patients. *Ann Surg Treat Res* 2014, 86(5):249-55.
19. Koivusalo A, Pakarinen M, Rintala R: Laparoscopic suture rectopexy in the treatment of persisting rectal prolapse in children: A preliminary report. *Surg Endosc* 2006; 20:960–3.
20. Portier G, Iovino F, Lazorthes F: Surgery for rectal prolapse: Orr-Loygue ventral rectopexy with

- limited dissection prevents postoperative-induced constipation without increasing recurrence. *Dis Colon Rectum* 2006; 49(8):1136-40.
21. Ruiz Carmona, Martín Arévalo J, Moro Valdezate D, et al: Sacral nerve stimulation for the treatment of severe faecal incontinence: Results after 10 years experience. *Cir Esp* 2014; 92(5):329-335.
 22. Bharucha AE: Outcome measures for fecal incontinence: anorectal structure and function. *Gastroenterology* 2004; 126(1): S90-8.
 23. Farouk R, Duthie GS, MacGregor AB, et al: Rectoanal inhibition and incontinence in patients with rectal prolapse. *Br J Surg* 1994; 81(5):743-6.
 24. Fahmy MA, Ezzelarab S: Outcome of submucosal injection of different sclerosing materials for rectal prolapse in children. *Pediatr Surg Int* 2004; 20 : 353 – 356.
 25. Kim DS, Tsang CB, Wong WD: Complete rectal prolapse: evolution of management and results. *Dis Col Rectum* 1999;42: 460 – 466.
 26. Jacobs LK, Lin YJ, Orkin BA: The best operation for rectal prolapse. *Surg Clin North Am* 1997; 77: 49 – 70.